

MATH 592 SPRING 2010
TOPICS IN ERGODIC THEORY

PROBLEM SET 4

1. Show that T is ergodic if and only if the only T -invariant measurable functions are the functions which are constant almost everywhere.
2. Show that the map $D : \mathbb{T} \rightarrow \mathbb{T}$ which sends $x \mapsto 2x$ is ergodic.
3. Give an example to show that it is possible for T and T^{-1} to be ergodic, without T^n being ergodic for all n .
4. Prove the following extension of Poincaré's recurrence theorem. If (X, \mathcal{X}, μ, T) is a measure-preserving system, and $f \in L^1(X, \mathcal{X}, \mu, T)$ is non-negative, then

$$\limsup_{n \rightarrow \infty} \int_X f T^n f \geq \left(\int_X f d\mu \right)^2.$$

5. Give an example to show that even in an ergodic system, the double average

$$\frac{1}{N} \sum_{n=0}^{N-1} f_1(T^n x) f_2(T^{2n} x)$$

does not necessarily converge to a constant function.

6. To become familiar with Fourier analysis on \mathbb{T} , prove at least two of the following statements:
 - (1) Define the N th Fejér kernel $K_N : \mathbb{T} \rightarrow \mathbb{R}$ by $K_N(\theta) = \frac{1}{2N+1} \sum_{n=-N}^N e^{2\pi i n \theta}$. Show that $f * K_N \rightarrow f$ uniformly if $f \in C(\mathbb{T})$.
 - (2) Show that $f * K_N \rightarrow f$ in L^1 if $f \in L^1(\mathbb{T})$.
 - (3) For $f \in L^1(\mathbb{T})$, write $f * K_N$ in terms of the Fourier coefficients $\widehat{f}(t)$.
 - (4) Show that if $f, g \in L^1(\mathbb{T})$ and if $\widehat{f}(t) = \widehat{g}(t)$ for all $t \in \mathbb{Z}$, then $f = g$ almost everywhere.
 - (5) For $f \in C^\infty(\mathbb{T})$, show that $\sum_t \widehat{f}(t) e^{2\pi i t \theta}$ converges uniformly to f .
 - (6) For $f \in L^2(\mathbb{T})$, show that the partial sums $\sum_{|t| \leq N} \widehat{f}(t) e^{2\pi i t \theta}$ converge to f in L^2 .

Please send any comments or corrections to jwolf137@math.rutgers.edu.